



Bilkent University  
Department of Mathematics

**Quiz # 8**  
Math 101-Section 09 Calculus I  
19 November 2015, Thursday



Instructor: Ali Sinan Sertöz


YOUR NAME:

**In this quiz you can use only pencils and erasers.**

*Show your work in detail, unless only an answer is required. Correct answer without proper explanation does not receive any partial credits.*

**Q-1)** Let  $g(x) = \int_0^x f(t) dt$ , where  $f$  is the function whose graph is shown.

- (a) At what values of  $x$  do the local maximum and local minimum values of  $g$  occur?
- (b) Where does  $g$  attain its absolute maximum value?
- (c) On what intervals is  $g$  concave downward?
- (d) Sketch the graph of  $g$ .



**Answer:** (a) By FTC1,  $g'(x) = f(x)$ . So  $g'(x) = f(x) = 0$  at  $x = 1, 3, 5, 7$ , and  $9$ .  $g$  has local maxima at  $x = 1$  and  $5$  (since  $f = g'$  changes from positive to negative there) and local minima at  $x = 3$  and  $7$ . There is no local maximum or minimum at  $x = 9$ , since  $f$  is not defined for  $x > 9$ .

(b) We can see from the graph that  $|\int_0^1 f dt| < |\int_1^3 f dt| < |\int_3^5 f dt| < |\int_5^7 f dt| < |\int_7^9 f dt|$ . So  $g(1) = |\int_0^1 f dt|$ ,  $g(5) = \int_0^5 f dt = g(1) - |\int_1^3 f dt| + |\int_3^5 f dt|$ , and  $g(9) = \int_0^9 f dt = g(5) - |\int_5^7 f dt| + |\int_7^9 f dt|$ . Thus,  $g(1) < g(5) < g(9)$ , and so the absolute maximum of  $g(x)$  occurs at  $x = 9$ .

(c)  $g$  is concave downward on those intervals where  $g'' < 0$ . But  $g'(x) = f(x)$ , so  $g''(x) = f'(x)$ , which is negative on (approximately)  $(\frac{1}{2}, 2)$ ,  $(4, 6)$  and  $(8, 9)$ . So  $g$  is concave downward on these intervals.

